

Risk Factors and Prevalence of Taeniasis among Displaced peoples in Ombadda Locality- Omdurman -Sudan

Jamal Yousef Lowaty Lowba^{1*}, Mahjoob Mohamed Ahmed¹, Fawzy Ahmed Idrees¹, Samwell Khalid Ahmed¹, Abualgasim Mohamed Anjool¹, Amir Mohamed Saleh Ali¹

¹ Omdurman Islamic University, Faculty of Medical Laboratory science.

Abstract

Background. Taeniasis is infection of human with intestinal cestode tape worm parasites belonged to the genus *Taenia*, species *saginata* and *solium*.

Objective: The study aimed to determine the risk factors and prevalence of taeniasis among displaced peoples district in Ombadda Omdurman state- Sudan.

Material and Methods: This is a cross sectional study was conducted in Ombadda and Basheer hospitals in Omdurman-Sudan from the period September to November 2021. A total of 150 stools samples were collected and examined using direct microscopy, formal ether concentration technique and Ziehl-Neelsen staining method for detecting *Taenia* Spp.

Results: A total of 150 stools samples collected and examined, 10 (6.3%) out of 150 faecal samples were positive and 140 (93.7%) were negative sample. Out of 10 positive samples 6 (60%) were males and 4(40%) were females their age groups range from 20 – 60 Years old, The Eggs stage of *Taenia saginata* and *Taenia solium* was identified by using direct microscopy and formal ether concentration techniques and differentiated by the Ziehl-Neelson staining method.

Key words: Taeniasis, risk factors of transmission, displaced peoples.

Corresponding author: Jamal Yousef Lowaty Lowba Tel: 0122075283, [jamal012275@gmail.com/](mailto:jamal012275@gmail.com)

Introduction:

Taeniasis is infection of human with intestinal tapeworms cestode parasites belonged to the genus *Taenia* and It is a significant zoonotic disease (6) Because it may cause cysticercosis in the tapeworm carriers, family members, and other close contacts (17). The most important human pathogen species in the genus *Taenia* are *Taenia solium* the pork tapeworm and *Taenia saginata* the beef tapeworm. The other species *Taenia asiatica* is found only in East Asia (9). *Taenia saginata* and *Taenia solium* are worldwide distributed, *taenia saginata* found in

the countries where cattle are raised a while *Taenia solium* is most common found in latin America, Southeast Asia and Africa practically prevalent in rural area where domestic pigs are found (3 ,19). Human infected with *Taenia saginata* and *Taenia solium* when they consumption infected beef meats or pigs' liver which were poorly cooked and eaten (8). These parasites have been complete indirect life cycle in two hosts, the definitive host was human which harbors the adult worm a while Porks and Cattles are intermediate host which was harbors the larval infective stage (15). Eggs or

gravid proglottids are passed with feces of human the eggs are highly infectious and can survive for days to months in different environmental sources (7). Cattles and pigs become infected by ingesting vegetation contaminated with eggs or gravid proglottids. *Taenia saginata* and *Taenia solium* consider to be like other several parasitic diseases cause mortality, *T. solium* cysticercosis is one of the most lethal parasitic diseases and is the most important foodborne Parasite (16). In terms of health and economic burden, *T. solium* has been reported as the first, and *Taenia saginata* as the nineteenth foodborne parasite at the global level [4]. *Taenia saginata* has a global distribution in Sudan than *T. solium* because it is considered one of the public health problem it is increasing mostly in the area where catles were arizing in community and if untreated causes economic burden and sever complication in human.

Material and Methods:

A cross sectional study conducted in Ombadda and Basheer hospitals in Omdurman State-Sudan from September to December 2021. A total of 150 faecal samples were collected in containers contain 10% formal saline as preservative reagents. were examined by microscopy and forma ether concentration as diagnostic methods, preserved faeses in formal saline were emulsified and homogenized, one drop of emulsified faecse suspensions placed on a slide microscope, covered with cover class, examined microscopy by wet preparation technique, using the 10 x objective and 40x objective of microscope lens for detecting segments and eggs of *Taenia* SPP in faeces. Using formal ether concentration method, 1 g of faeces

emulsified in 4 ml of 10% forma water in test tubes, 3-4 ml of formal water added further in test tubes mixed well, emulsified faeces sieved in beaker, 2-3 ml of diethyl ether added to the suspension and mixed in the test tubes, homogenate suspension was centrifuged at 750-1000 g for 1 minute and the sediment was examined microscopically by using the 10 X objective and 40 x to identify eggs and segment of *Taenia* Spp. The Eggs stage of *Taenia saginata* and *Taenia solium* was detected, identified and differentiated by the Ziehl-Neelson staining method.

Results

A total of 150 faecal samples collected from displaced peoples in Ombada hospitals Omdurman - state examined. 10 (6.3%) out of 150 faecal samples were positive and 140(93.7%) were negative samples for *Taenia* Spp (Table1). The 10 positive samples that were seen in faecal samples *Taenia saginata* was9 (90%) and *Taenia solium* 1(10%) the eggs stag of *Taenia* Spp was identified using direct microscopy and formal ether concentration as diagnostic methods and differentiated by the Ziehl-Neelson staining method (Table3). Among these10 positive samples 6(60 %) were found to occur in males' and 4 (40 %) were found in female's patients the difference was found to be statistically in significant (*P. Value*=0.454) as shown in (Table2) .All patients came from displaced area their age groups range from 20 – 60 Years old and The age groups 31-40 to51-60 Years old were most commonly infected by *Taenia* SPP this differences was found to be statistically in significant (*P. Value*=0.425) (

Table 2) The Infection rate decreased in Patients their jobs 1(10%) were students and 2(20%) were non workers this differences was found to be statistically in significant (P.Value=0.566)in (Table 2) . Patients jobs 5(50%) Slaughters and 2(20%) farmers were more susceptible to Taenia Spp infection in both sexes than other occupied else where the differences was found to be statistically significant (P. Value 0.001) (Table 2). High infection rate occur in Patients had past history of

eating under cooked meat of beef 9(90%) and eating under cooked meats of pork 1(10%) there was correlation association between Taenia SPP infection and peoples has cultural practices eating under cooked meat of beef and meat of pork the differences was found to be statistically significant (P. Value 0.001) (Table 4.). infection rate increased in patients not using toilet for defecation and decreased in patients using toilet for stool disposable (Table 4).

Table (1): Prevalence of Taeniasis among displaced peoples in Ombadda and Basheer hospitals Omdurman state.

No Examined	Taenia SPP		Total Number
	Positive%	Negative%	
150	10(6.7%)	140(93.3%)	150

Table (2): Demographic characteristics of 10 patients with Taeniasis in Omdurman state.

Variables	Taeniasis SPP		Total No	P-Value
Gender		T.saginat %	T.solium %	
	Males	5(60%)	1(10%)	10 (100%)
	Females	4 (40%)	0 (0%)	
Ages	20-30	2(20%)	0(0%)	2(20% (
	31-40	2(20%)	1 (10 %)	3(30%)
	51-60	5 (50%)	0(0%(	5(50%)
	Total	9(90%)	1(10%)	10(100%)
Occupations	Student	1(10%)	0 (0%)	1(10%)
	Non workers	1(10%)	1(10%)	1(10%)
	Farmers	2(20%)	0 (0%)	2(20%)
	Slaughters	5 (50%)	1(10%)	1(10%)
	Total	9(90%)	1(10%)	10(100%)

Tables (3): Morphological differences between *T. saginata* and *T. solium* eggs using the Ziehl-Neelson staining method in 10 patients in Ombada hospital.

Total of feacal examined	Taeniasis positive	Ziehl-Neelson		Total No
		T.saginat	T.solium	
150	10	9(90%)	1 (10%)	10(100%)

Table (4) prevalence of Taeniasis and associated risk factors in displaced peoples

History of eating raw meats	T.saginata	T.solium	Total No	P.Value
Pork raw consumption +ve (1)	0 (0 %)	1(10%)	9(90%)	
Beef raw consumption +ve	9(90%)	0(0%)	1(10%)	0.001
Total	9(90%)(	1 (10%)	10(100%	
Method of stool disposable				
Open defecation yes (8)	8 (80%)	0(0%)		
Using Toilet yes (2)	1(10%)	1(10%)		0.001
Total	9(90%)	1(10%)	10(100%)	

Discussions:

The results obtained in the present study indicated that the overall infection rate of Taeniasis among the displace people was 6.4% this percentage results considered to be higher rate of infection when compared by previous study obtained by(Ghada et al., 2016) (10) found that(1.6%) in Sudan and higher than percentage results by(McCleery et al.,2015) (13) reported a low prevalence rate(2.9%) of taeniasis among refugees living on the Thai–Myanmar border in Thailand and were extremely lower when compared by previous results reported by(wongsaroi et al., 2014)(20) in Korea reported that 18.2% of Taeniasis among Korean people probably due to consumption raw meats or under cooked meats of beefs and porks which was containing infective

stage of Taeniais . Our results in the present study showed that *Taenia saginata* infection rate was 9(90%) and *Taenia solium* infection rate was 1(10%) this indicate that *Taenia saginata* was more prevalent in displace people than *Taenia solium* due to Muslim religion in the area and population raises cattle's houses, for food or for breeding. which was risk factor for taeniasis this results obtained when compared by previous study by (Anantaphruti et al., 2007) (2) in Thailand showed significant differences. When we analyzed the taeniasis infection in the gender, the prevalence rate was highest in males 6(60%) and lower in females 4(40%) the middle-aged groups ranging from 20–60 years old to 31–40 years old and 51-60 Years old were most commonly infected by *Taenia saginata* this significant differences of

a higher infection in males agreement with previous study by (Ng-Nguyen et al., 2018) (14) reported in highlands of Vietnam where it was noted that males has history of consumption raw meats and undercooked meats of beefs and pork's as cultural practices in dis place area .The results obtained showed that there are no Taeniasis infection reported in children age group(< 20 Years old)this may be due to fact that the samples size of stools was collected in sufficient to declared that children are not in risk of infection by Taeniasis when compare this results with previous studies showed that highly significant differences has been reported by (Lit et al.,2019) (11) in China ,(Madinga et al.,2012)(12) in Chan go and (Xud-M et al., 2010)(21) in Philippines reported that the highly Taeniasis infection found in children age group(< 20 Years old) .In the present study the Infection rate of Taeniasis carriers decreased in Patients their jobs 1(10%) were students and 2(20%) were non workers this results agree with the previous results reported by (Openshaw et al., 2018) (1) in western Sichuan, People's Republic of China and were predominantly occur in Patients their jobs 5(50%) Slaughters and 2(20%) farmers infection in both sexes than other occupied elsewhere this differences declared that the displace peoples a history of consuming raw or undercooked raw pork. , beef raw meats , working in corn fields and by cooking at their work places which was the most predominant risks factor for prevalence of Taeniasis similar results reported by (Swastika et al.,2017) (18) in Karangasem Villages

Indonesia .The present study showed that infection rate increased in patients has history of passing stools outside their houses not using toilet for defecation and decreased in patients using toilet for stool disposable which pre-oral infection by Taenia eggs These are important risk factors of taeniasis when compared by previous results reported by (Ito et al., 2020) (23) showed similarity.

Conclusion: Taeniasis is the major public health problem mostly associated with those have poor hygiene habit, lack of meat inspection, bush defecation and poor waste disposable which has been considered risk factors for Taeniasis, slaughtering practices have contributed to the establishment and transmission of disease, Taenia saginata is more common and prevalent indisplaced people than Taenia solium, consumption under cooked Beef meats and pork meats has been hazardous and risk factors facilitated transmission of Taeniasis.The results obtained showed that the Ziehl-Neelson staining method used more effective in parasites differencation than other techniques used

Conflict of interest: there are no conflict of interest.

Financial: this study has not received financial support.

Acknowledgment: I would to appreciate laboratory staph of Ombada hospitals and Basheer hospitals in Omdurman locality -Sudan.

Reference:

1. Openshaw JJ, Medina A, Felt SA, Li T, Huan Z, Rozelle S,Luby SP. 2018. Prevalence and risk

- factors for *Taenia solium* cysticercosis in school-aged children: A school based study in western Sichuan, People's Republic of China. *PLoS Neglected Tropical Diseases*, 12(5), e0006465.
2. Anantaphruti MT, Yamasaki H, Nakao M, Waikagul J, Watthanakulpanich D, Nuamtanong S, Maipanich W, Pubampen S, Sanguankiat S, Muennoo C, Nakaya K, Sato MO, Sako Y, Okamoto M, Ito A. 2007. Sympatric occurrence of *Taenia solium*, *T. saginata*, and *T. asiatica*, Thailand. *Emerging Infectious Diseases*, 13(9),
 3. Kaewpitoon SJ, Loyd RA, Kaewpitoon N. 2015. A crosssectional survey of intestinal helminthiasis in rural communities of Nakhon Ratchasima province, Thailand. *Journal of the Medical Association of Thailand*, 98(suppl 4), 27–32.
 4. WHO. (2014). Multicriteria-based ranking for risk management of food-borne parasites: report of a Joint FAO: FAO, World Health Organization pp.
 5. Yamasaki H, Allan JC, Sato MO, Nakao M, Sako Y, Nakaya K, Qiu D, Mamuti W, Craig PS, Ito A. 2004. DNA differential diagnosis of taeniasis and cysticercosis by multiplex PCR. *Journal of Clinical Microbiology*, 42(2), 548–553.
 6. Acha PN, Szyfres B. (2003). Zoonoses and communicable diseases common to man and animals. Volume 3. Parasitoses. 3rd ed. Washington DC: PAHO; 2003. Scientific.
 7. Adewole SO (2010) Prevalence and Epidemiological Factors Influencing the Transmission of Human Taeniasis in Ekiti State, Nigeria. *International Journal of Academic Research* 2: 90-93.
 8. Garcia HH, Gonzalez AE, Evans CA, Gilman RH, Peru CWGi. 2003. *Taenia solium* cysticercosis. *Lancet*, 362(9383), 547–556.
 9. Garcia, Oscar H, Del Brutto, Hector H. (2014). *Taenia solium* biological characteristics and life cycle. *cysticercosis of human nervous system*. Berlin. P.11-12.
 10. Ghada Hassan, Shawgi Hassan, Atif Elamin (2016) Meat-borne parasite a health hazard concern in Sudan. P.105-106.
 11. Li T, Chen X, Wang H, Openshaw JJ, Zhong B, Felt SA, Ito A, Luby SP. 2019. High prevalence of taeniasis and *Taenia solium* cysticercosis in children in western Sichuan, China. *Acta Tropica*, 199, 105133.
 12. Madinga J, Kanobana K, Lukanu P, Abatih E, Baloji S, Linsuke S, Praet N, Kapinga S, Polman K, Lutumba P. 2017. Geospatial and age-related patterns of *Taenia solium* taeniasis in the rural health zone of Kimpese, Democratic Republic of Congo. *Acta Tropica*, 165, 100–109.
 13. McCleery EJ, Patchanee P, Pongsopawijit P, Chailangkarn S, Tiwananthagorn S, Jongchansittoe P, Dantrakool A, Morakote N, Phyu H, Wilkins PP. 2015. Taeniasis among refugees living on Thailand-Myanmar border, 2012. *Emerging Infectious Diseases*, 21(10), 1824.
 14. Ng-Nguyen D, Stevenson MA, Breen K, Van Phan T, Nguyen V-AT, Van Vo T, Traub RJ. 2018. The epidemiology of *Taenia* spp. infection and *Taenia solium* cysticerci exposure in humans in the Central Highlands of Vietnam. *BMC Infectious Diseases*, 18(1), 1–9.

15. O'Neal S, Noh J, Wilkins P, Keene W, Lambert W, et al. (2011) *Taenia solium* Tapeworm Infection, Oregon, 2006-2009. *Emerg Infect Dis* 17: 1030-1036.
16. Prasad KN, Prasad A, Gupta RK, Pandey CM, Singh U (2007) Prevalence and associated risk factors of *Taenia solium* Taeniasis in a rural pig farming community of north India. *Trans R Soc Trop Med Hyg* 101: 1241-1247.
17. Schantz PM, Cruz M, Sarti E, Pawlowski ZJ. 1993. Potential eradicability of taeniasis and cysticercosis. *Bulletin of the Pan American Health Organization*, 27(4), 397–403.
18. Swastika K, Wandra T, Dharmawan NS, Sudarmaja IM, SaragihJM, Diarthini LPE, Ariwati L, Damayanti PAA, Laksemi DAAS, Kapti N, Sutisna P, Yanagida T, Ito A. 2017. Taeniasis caused by *Taenia saginata* in Gianyar town and *Taenia solium* in Karangasem villages of Bali, Indonesia, 2011–2016: How to detect tapeworm carriers, anamnesis or microscopy? *Acta Tropica*, 174, 19–23.
19. Wandra T, Swastika K, Dharmawan NS, Purba IE, Sudarmaja IM, Yoshida T, Sako Y, Okamoto M, Diarthini NLPE, Laksemi DAAS, Yanagida T, Nakao M, Ito A. 2015. The present situation and towards the prevention and control of neurocysticercosis on the tropical island, Bali. Indonesia. *Parasites & Vectors*, 8(1), 1–11.
20. White AC Jr. 2000. Neurocysticercosis: updates on epidemiology, pathogenesis, diagnosis, and management. *Annual Review of Medicine*, 51(1), 187–206.
21. Wongsaroj T, Nithikathkul C, Rojkitikul W, Nakaia W, Royal L, Rammasut P. 2014. National survey of helminthiasis in Thailand. *Asian Biomedicine*, 8(6), 779–783.
22. Xu J-M, Acosta LP, Hou M, Manalo DL, Jiz M, Jarilla B, PabloAO, Ovleda RM, Langdon G, McGarvey ST, Kurtis JD, Friedman JF, Wu H-W. 2010. Seroprevalence of cysticercosis in children and young adults living in a helminth endemic community in Leyte, the Philippines. *Journal of Tropical Medicine*, 2010, 603174.
23. Zoli A, Shey-Njila O, Assana E, Nguekam JP, Dorny P, et al. (2003) Regional status, epidemiology and impact of *Taenia solium* cysticercosis in Western and Central Africa. *Acta Tropica* 87: 35-42.
24. Ito A, Saito M, Donadeu M, Lightowlers MW. 2020. KozenYoshino's experimental infections with *Taenia solium* tapeworms: An experiment never to be repeated. *Acta Tropica*, 205, 105378.